

Two-nucleon separation energies within the proxy-SU(3) model

S. Sarantopoulou, Andriana Martinou, Dennis Bonatsos

Institute of Nuclear and Particle Physics, National Centre for Scientific Research Demokritos, GR-15310 Aghia Paraskevi, Attiki, Greece

Abstract

Proxy-SU(3) is an approximate symmetry appearing in heavy deformed nuclei [1,2]. In SDANCA-2017 the foundations of proxy-SU(3) [3], its parameter-free predictions for the collective deformation parameters β and γ [4,5], as well as for $B(E2)$ ratios [5], have been discussed and its usefulness in explaining the dominance of prolate over oblate shapes in the ground states of even-even nuclei [6] and the point of the prolate to oblate shape transition in the rare earths region [6] has been demonstrated. In the present contribution, preliminary calculations for two-nucleon separation energies within this framework will be discussed.

References

- [1] D. Bonatsos, I.E. Assimakis, N. Minkov, A. Martinou, R.B. Cakirli, R.F. Casten, and K. Blaum, *Phys. Rev. C* **95** (2017) 064325.
- [2] D. Bonatsos, I.E. Assimakis, N. Minkov, A. Martinou, S. Sarantopoulou, R.B. Cakirli, R.F. Casten, and K. Blaum, *Phys. Rev. C* **95** (2017) 064326.
- [3] I.E. Assimakis, D. Bonatsos, N. Minkov, A. Martinou, R.B. Cakirli, R.F. Casten, and K. Blaum, *Bulg. J. Phys.* **44** (2017) 398-406.
- [4] D. Bonatsos, I.E. Assimakis, N. Minkov, A. Martinou, S.K. Peroulis, S. Sarantopoulou, R.B. Cakirli, R.F. Casten, and K. Blaum, *Bulg. J. Phys.* **44** (2017) 385-397.
- [5] A. Martinou, D. Bonatsos, I.E. Assimakis, N. Minkov, S. Sarantopoulou, R.B. Cakirli, R.F. Casten, and K. Blaum, *Bulg. J. Phys.* **44** (2017) 407-416.
- [6] S. Sarantopoulou, D. Bonatsos, I.E. Assimakis, N. Minkov, A. Martinou, R.B. Cakirli, R.F. Casten, and K. Blaum, *Bulg. J. Phys.* **44** (2017) 417-426.